

PHC Vital Signs Profile: Indicator Description Sheets

ARGENTINA

List of Indicators

Context	3
1. GDP per capita (PPP current international \$)	3
2. Population living in poverty (Under \$1.90 int'l dollars / day).....	4
3. Government health spending as percentage of GDP	5
Outcomes.....	6
4. Life expectancy at birth (years)	6
5. Maternal mortality ratio (per 100,000 live births)	7
6. Neonatal mortality rate (per 1,000 live births).....	8
7. Premature noncommunicable disease (NCD) mortality (probability)	9
8. Causes of death	10
Financing.....	11
Capacity	11
Performance	11
<i>Access</i>	<i>11</i>
<i>Quality.....</i>	<i>12</i>
Comprehensiveness	12
9. Average availability of RMNCH services	12
10. Average availability of NCD services.....	13
Continuity.....	14
11. DTP3 dropout rate.....	14
12. Treatment success rate for new TB cases.....	15
Person-Centeredness	15
Provider Availability	15
Provider Competence.....	16
13. Early antenatal care.....	16
14. Adherence to guidelines for child health	17
15. Adherence to guidelines for adolescent health	18
Safety	18
<i>Service Coverage.....</i>	<i>19</i>
RMNCH.....	19
16. Demand for family planning satisfied with modern methods	19
17. Antenatal care coverage (4+ visits)	20
18. Coverage of DTP3 immunization	21
19. Care-seeking for suspected child pneumonia	22
Infectious Diseases.....	23
20. TB cases detected and treated with success	23
21. People living with HIV receiving anti-retroviral treatment.....	24
22. Children under 5 with diarrhea receiving ORS.....	25
Noncommunicable Diseases	26
1. Prevalence of raised blood pressure (age-standardized estimate).....	27

Equity.....	27
2. Coverage of RMNCH services, by wealth quintile.....	27
3. Under-five mortality rate, by wealth quintile of provinces	28

Context

Context indicators cover important contextual details about a country, including GDP per capita, the proportion of the population living in poverty, and government spending on health.

1. GDP per capita (PPP current international \$)

Full Name of Indicator	GDP per capita, PPP (current international dollars)
Short name of indicator	GDP per capita (\$PPP international dollars)
Description	Gross domestic product per capita converted to international dollars using purchasing power parity rates. An international dollar has the same purchasing power over GDP as the U.S. dollar has in the United States. Data are in current international dollars.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Context
Construction	<p><i>Numerator:</i> GDP is the sum of gross value added by all resident producers in the economy plus any product taxes and minus any subsidies not included in the value of the products. It is calculated without making deductions for depreciation of fabricated assets or for depletion and degradation of natural resources.</p> <p><i>Denominator:</i> Total population</p>
Rationale	GDP per capita is an important contextual indicator that provides information about the average annual income of country residents.
Data Source & Year	World Development Indicators (World Bank), 2017.
Limitations	<p>GDP as a measure has some limitations including: (1) it doesn't capture non-market production; (2) it doesn't capture underground or non-official economies; (3) it doesn't measure the possible negative effects (e.g. on quality of life or environment of the production captured in the measure; and (4) trending can be difficult due changes in the quality of products and the inclusion of new goods.</p> <p>Additionally, GDP estimates can vary greatly depending on the basket of goods captures and the currency used for reporting. There may be differences in national accounting and demographic reporting procedures and practices between countries.</p>
VSP Methodology	N/A

2. Population living in poverty (Under \$1.90 int'l dollars / day)

Full Name of Indicator	Proportion of population below international poverty line of \$1.90 per day (2011 PPP)
Short name of indicator	% Living in poverty
Description	Percentage of the population living in poverty, defined as living on less than \$1.90 international dollars per day. An international dollar has the same purchasing power over GDP as the U.S. dollar has in the United States. Data are in constant 2011 international dollars.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Context
Construction	<i>Numerator:</i> Total population living on less than \$1.90 international dollars per day <i>Denominator:</i> Total population
Rationale	Populations living in poverty may face greater barriers to health services access and utilization.
Data Source & Year	World Development Indicators (World Bank), 2016. Data are based on primary household survey data obtained from government statistical agencies and World Bank country departments.
Limitations	The timeliness, frequency, quality, and comparability of household surveys may be poor, particularly in the poorest countries. The availability and quality of poverty monitoring data remains low in small states, countries with fragile situations, and low-income countries and even some middle-income countries.
VSP Methodology	N/A

3. Government health spending as percentage of GDP

Full Name of Indicator	Domestic General Government Health Expenditure as % of Gross Domestic Product (GDP)
Short name of indicator	Government health spending as % of GDP
Description	<p>Domestic General Government Health Expenditure as % of GDP measures current government expenditure on health, from domestic sources, relative to the country's GDP. Domestic General Government Health Expenditure tracks expenditure by all public and compulsory sources for health, exclusively from domestic revenue.</p> <p>The numerator refers to health care goods and services used or consumed during a year. Note that capital investments are excluded.</p>
Comparability	Comparable/Standard indicator
VSP Domain and Sub-Domain	Financing
Construction	<p><i>Numerator:</i> Domestic General Government Health Expenditure</p> <p><i>Denominator:</i> Gross Domestic Product (GDP)</p>
Rationale	Contributes to understanding overall government expenditure on health in relation to the size of the national economy.
Data Source & Year	WHO Global Health Expenditure Database, 2015.
Limitations	In this case, the indicator value presented differs from country data sources due to the adoption of methods to enhance international comparability. The value as calculated from national data is 7.1% (Hacienda, Gasto Público Consolidado en Salud (nación, provincias y municipios), 2015).
VSP Methodology	N/A

Outcomes

Outcomes focus on the health status of the population, including life expectancy, mortality, and causes of death.

4. Life expectancy at birth (years)

Full Name of Indicator	Life expectancy at birth (years)
Short name of indicator	Life expectancy
Description	The average number of years that a newborn could expect to live if he or she were to pass through life exposed to the sex- and age-specific death rates prevailing at the time of his or her birth, for a specific year, in a given country, territory or geographical area.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Outcomes
Construction	Life expectancy at birth is derived from life tables and is based on sex- and age-specific death rates. United Nations values for life expectancy at birth correspond to mid-year estimates, consistent with the corresponding United Nations fertility medium-variant quinquennial population projections. Procedures used to estimate WHO life tables for Member States vary depending on the data available to assess child and adult mortality.
Rationale	Life expectancy at birth is one of the key measures of a population's health and is a reflection of the overall mortality level and pattern across all age groups within the population.
Data Source & Year	Global Health Observatory (GHO), 2016. Data on maternal mortality and other relevant variables are obtained through databases maintained by WHO, UNPD, UNICEF, and the World Bank. Data available from countries vary in terms of the source and methods. Given the variability of the sources of data, different methods are used for each data source in order to arrive at country estimates that are comparable and permit regional and global aggregation.
Limitations	The lack of complete and reliable mortality data, especially for low income countries and particularly on mortality among adults and the elderly, necessitates the application of modelling (based on data from other populations) to estimate life expectancy. This may lead to minor differences compared with official life tables prepared by Member States.
VSP Methodology	N/A

5. Maternal mortality ratio (per 100,000 live births)

Full Name of Indicator	Maternal mortality ratio (per 100,000 live births)
Short name of indicator	Maternal mortality ratio
Description	The annual number of female deaths from any cause related to or aggravated by pregnancy or its management (excluding accidental or incidental causes) during pregnancy and childbirth or within 42 days of termination of pregnancy, irrespective of the duration and site of the pregnancy, expressed per 100,000 live births, for a specified time period.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Outcomes
Construction	<i>Numerator:</i> Number of maternal deaths <i>Denominator:</i> Number of live births (expressed per 100,000 live births)
Rationale	Complications during pregnancy and childbirth are a leading cause of death and disability among women of reproductive age in developing countries. The maternal mortality ratio represents the obstetric risk associated with each pregnancy and monitors deaths related to pregnancy and childbirth. It reflects the capacity of the health system to provide effective health care in preventing and addressing the complications occurring during pregnancy and childbirth that can result in maternal death.
Data Source & Year	Global Health Observatory (GHO), 2015. Data on maternal mortality and other relevant variables are obtained through databases maintained by WHO, UNPD, UNICEF, and the World Bank. Data available from countries vary in terms of the source and methods. Given the variability of the sources of data, different methods are used for each data source in order to arrive at country estimates that are comparable and permit regional and global aggregation.
Limitations	Vital registration and health information systems in most developing countries are weak and thus cannot provide an accurate assessment of maternal mortality. Even estimates derived from complete vital registration systems, such as those in developed countries, suffer from misclassification and underreporting of maternal deaths.
VSP Methodology	N/A

6. Neonatal mortality rate (per 1,000 live births)

Full Name of Indicator	Neonatal mortality rate (probability of dying within the first 28 days of life per 1,000 live births)
Short name of indicator	Neonatal mortality rate
Description	The neonatal mortality rate is the probability of a newborn dying before reaching 28 days of age, expressed per 1,000 live births.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Outcomes
Construction	Numerator: Number of deaths of neonates at ages 0-28 days Denominator: Number of live births for a specified year (expressed per 1,000 live births)
Rationale	Mortality during the neonatal period accounts for a large proportion of child deaths and is considered to be a useful indicator of maternal and newborn neonatal health care. Neonatal mortality rate is a Sustainable Development Goal Indicator for monitoring child health.
Data Source & Year	UN IGME, 2016. The Inter-agency Group for Child Mortality of Estimation, which includes representatives from UNICEF, WHO, the World Bank and the United Nations Population Division, produces trends of neonatal mortality with standardized methodology by group of countries depending on the type and quality of source of data available. These neonatal rates are estimates, derived from the estimated UN IGME neonatal rate and infant population from World Population Prospects to calculate the live births; hence they are not necessarily the same as the official national statistics.
Limitations	The reliability of estimates of neonatal mortality depends on the accuracy and completeness of reporting and recording of births and deaths. Underreporting and misclassification are common.
VSP Methodology	N/A

7. Premature noncommunicable disease (NCD) mortality (probability)

Full Name of Indicator	Mortality between ages 30 and 70 years from cardiovascular diseases, cancer, diabetes or chronic respiratory diseases (probability)
Short name of indicator	Premature NCD mortality
Description	Probability of dying between the ages of 30 and 70 years from non-communicable diseases, defined as the percent of 30-year-old-people who would die before their 70th birthday from cardiovascular disease, cancer, diabetes, or chronic respiratory disease, assuming that s/he would experience current mortality rates at every age and s/he would not die from any other cause of death.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Outcomes
Construction	<i>Numerator:</i> Number of deaths between ages 30 to 70 years from cardiovascular disease, cancer, diabetes, or chronic respiratory disease in a synthetic life table population. <i>Denominator:</i> Population at exact age 30 in the synthetic life table population.
Rationale	Non-communicable diseases account for an increasing proportion of morbidity and mortality in many countries. Prevention, diagnosis, and treatment of these diseases to avoid premature mortality are a critical part of primary health care.
Data Source & Year	Global Health Observatory (GHO), 2016. Data are derived from re-analysis of Demographic and Health Surveys (DHS) micro-data, which are publicly available using the standard indicator definitions as published in DHS documentation.
Limitations	The reliability of estimates depends on the accuracy and completeness of reporting and recording of births and deaths. Underreporting and misclassification are common.
VSP Methodology	N/A

8. Causes of death

Full Name of Indicator	Cause-specific mortality
Short name of indicator	Causes of death
Description	Causes of death disaggregated by percentage attributable to non-communicable diseases (NCDs), injuries, and communicable and other conditions (including maternal, perinatal, and nutritional conditions).
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Outcomes
Construction	<i>Numerator:</i> Total number of deaths by cause in a given year <i>Denominator:</i> Total number of deaths in a given year
Rationale	Cause-of-death statistics allow governments to determine priorities for public health actions, such as increasing health spending in areas to which high mortality is attributed.
Data Source & Year	Global Health Observatory (GHO), 2016. Data are derived from re-analysis of Demographic and Health Surveys (DHS) micro-data, which are publicly available using the standard indicator definitions as published in DHS documentation.
Limitations	The reliability of estimates depends on the accuracy and completeness of reporting and recording of births and deaths. Underreporting and misclassification are common.
VSP Methodology	N/A

Financing

Financing includes measurements of per capita expenditures on Primary Health Care (PHC), share of health expenditure allocated to PHC, and health expenditures as percent of GDP.

No recent indicator available from international or national data sources.

Capacity

The PHC Progression Model uses mixed methods to assess foundational capacities of PHC. Scores for the Capacity domain are color-coded green (high, 4), yellow (medium, 2 or 3), or red (low, 1). Details for each of the measures can be found in PHC Progression Model Assessment Rubrics: DRAFT V1.1 <<https://improvingphc.org/phc-progression-model-assessment-rubrics-draft-v11>>.

Performance

The Performance domain includes measures of access, quality, and service coverage.

Where comparable data are available, scores for the Performance domain are color-coded green (good, 80+), yellow (medium, 60-79), or red (poor, <60). Scores based on data from non-comparable sources are colored gray.

Access

Access includes measurements of financial barriers and geographic hardship due to distance.

No recent indicator available from international or national data sources.

Quality

Quality of care measures are focused on principles that are proven to impact the quality of PHC service delivery at the point of care. These include comprehensiveness of care, continuity of care, person-centeredness, availability and competence of providers, and safety practices.

Comprehensiveness

9. Average availability of RMNCH services

Full Name of Indicator	Percentage of facilities offering services for the diagnosis and/or management of maternal and child health
Short name of indicator	Average availability of RMNCH services
Description	Percentage of facilities offering services for the diagnosis and/or treatment of maternal and child health, including antenatal care, services for children under 5 years old, family planning and vaccinations.
Comparability	Non-comparable / Country-specific proxy indicator for “Average availability of 5 tracer RMNCH services”
VSP Domain and Sub-Domain	Performance / Quality / Comprehensiveness
Construction	<i>Numerator:</i> Number of public facilities at the first level of care that report offering services for the diagnosis and treatment of maternal and child health, as per agreement <i>Denominator:</i> Number of public facilities at the first level of care that can report to the Program
Rationale	Maternal and child health services are a critical part of primary health care and must be widely available throughout all facilities in order to support maternal and child health.
Data Source & Year	Bases administrativas de la Secretaría de Gobierno de Salud de la Nación, 2018. In the numerator, the indicator considers facilities that have reported prenatal care services, services for children under 5 years old, family planning and vaccines, performed during the last 12 months. In the denominator, facilities that have the capacity to report services to the health system are included, that is, they have an agreement with the SUMAR Program. These centers represent 83% of the total establishments.
Limitations	While this indicator provides information on the extent to which maternal and child health services are offered, it does not fully assess the readiness of facilities to provide care that follows evidence-based practices, or the quality of the care itself. Use of an indicator reflecting questions related to assessing service readiness would provide additional information. This indicator does not include data from the private sector as it only considers those public facilities included in the Programa SUMAR.
VSP Methodology	N/A

10. Average availability of NCD services

Full Name of Indicator	Percentage of facilities offering services for the diagnosis and/or management of noncommunicable disease
Short name of indicator	Average availability of NCD services
Description	Proportion of facilities providing non-combinable services for the diagnosis and/or management of diabetes, cardiovascular diseases, chronic respiratory diseases, colon cancer, uterine cancer, overweight and obesity, smoking and chronic kidney disease.
Comparability	Non-comparable / Country-specific proxy indicator for “Average availability of diagnosis and management of 3 tracer NCDs”
VSP Domain and Sub-Domain	Performance / Quality / Comprehensiveness
Construction	<i>Numerator:</i> Number of facilities at the first level of care offering services for the diagnosis and treatment of NCDs <i>Denominator:</i> Number of facilities at the first level of care that can report to the Program
Rationale	Non-communicable diseases account for an increasing proportion of morbidity and mortality in many countries. Diagnosis and treatment of these diseases are a critical part of primary health care and must be widely available throughout all facilities in order to support the health of the population.
Data Source & Year	Bases administrativas de la Secretaría de Gobierno de Salud de la Nación, 2018. In the numerator, the indicator considers facilities that have reported diagnosis and / or treatment services for diabetes, cardiovascular diseases, chronic respiratory diseases, colon cancer, uterine cancer, overweight and obesity, smoking and chronic kidney disease, performed during the last 12 months. In the denominator, the indicator considers facilities that have the capacity to report services to the health system are included, that is, they have an agreement with the SUMAR Program. These centers represent 83% of the total facilities.
Limitations	While this indicator provides information on the extent to which NCD services are offered, it does not fully assess the readiness of facilities to provide care that follows evidence-based practices, or the quality of the care itself. Use of an indicator reflecting questions related to assessing service readiness would provide additional information. This indicator does not include data from the private sector as it only considers those public facilities included in the Programa SUMAR.
VSP Methodology	N/A

Continuity

11. DTP3 dropout rate

Full Name of Indicator	Dropout rate between 1 st and 3 rd DTP vaccination
Short name of indicator	DTP3 dropout rate
Description	Diphtheria-tetanus-pertussis (DTP) dropout rate is the percent of children who do not receive the full three doses of DTP vaccination after receiving the initial dose.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Quality / Continuity
Construction	<i>Numerator:</i> [DTP1 Immunization Coverage - DTP3 Immunization Coverage] <i>Denominator:</i> DTP1 Immunization Coverage
Rationale	Immunization is an essential component for reducing under-five mortality. Immunization coverage estimates are used to monitor coverage of immunization services and to guide disease eradication and elimination efforts. Measuring the gap between DTP1 and DTP3 reflects continuity within a health system, including the system's ability to capture and follow up with patients.
Data Source & Year	WHO/UNICEF, 2017. The WHO and UNICEF regularly report and release updated immunization coverage data related to the Global Vaccine Action Plan.
Limitations	Given the prevalence of global support for immunization efforts, a high coverage rate of DTP3 immunization may be reflective of strong support from vertical programming in some countries. As such, DTP3 coverage alone is not necessarily a proxy for primary care health system performance.
VSP Methodology	For calculation of summary scores in the VSP, this variable was transformed by subtracting the value from 100.

12. Treatment success rate for new TB cases

Full Name of Indicator	Tuberculosis cases detected and treated with success
Short name of indicator	Treatment success rate for new TB cases
Description	Percentage of tuberculosis (TB) cases successfully treated (cured plus treatment completed) among TB cases notified to national health authorities during a specified period, usually one year.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Quality / Continuity
Construction	<i>Numerator:</i> Number of TB cases registered in a specified time period that were successfully treated with or without bacteriological evidence of success <i>Denominator:</i> Total number of TB cases registered in the same period
Rationale	Treatment success is an indicator of the performance of national TB programs. It also serves as a proxy for a number of aspects of successful service delivery within a health system, including diagnostic and treatment accuracy and the system's ability to capture and follow up with patients over time.
Data Source & Year	WHO TB Programme, 2015. Preferred data sources include patient record and surveillance systems.
Limitations	This indicator measures only public-sector TB programs and does not include results from private-sector treatment programs or facilities. Therefore, in countries with strong private-sector TB programs, these results do not reflect the totality of the TB treatment success rate. Further, this indicator does not capture the system's ability to identify new TB patients. As a result, a country could perform well on this indicator, but poorly on the identification of new TB cases.
VSP Methodology	N/A

Person-Centeredness

No recent indicator available from international or national data sources.

Provider Availability

No recent indicator available from international or national data sources.

Provider Competence

13. Early antenatal care

Full Name of Indicator	Proportion of pregnant women with at least one check-up before the 13 th week of pregnancy
Short name of indicator	Early antenatal care
Description	Proportion of pregnant women with exclusive public coverage with at least one prenatal check-up before week 13 of gestation.
Comparability	Non-comparable / Country-specific proxy indicator for “Antenatal care quality score”
VSP Domain and Sub-Domain	Performance / Quality / Provider Competence
Construction	<p><i>Numerator:</i> Number of pregnant women with exclusive public coverage with at least one prenatal check-up before week 13 of gestation.</p> <p><i>Denominator:</i> Estimated number of pregnant women with exclusive public coverage.</p> <p>For the purpose of measuring the indicator, prenatal control is intended as the contact with the health system for diagnosis and / or monitoring of pregnancy with reported gestational age.</p>
Rationale	Argentina, following international recommendations, has identified the provision of early antenatal care as a key factor to capture the population of pregnant women and to ensure appropriate care during pregnancy can be delivered timely and according to guidelines.
Data Source & Year	Administrative data from the Secretaria de Gobierno de Salud, Ministerio de Salud y Desarrollo Social, 2018. For the denominator, data sources are CENSO (2010), EPH y EAHU.
Limitations	Data for this indicator comes from the Programa SUMAR, which provides insurance coverage to selected populations, and covers only public sector facilities.
VSP Methodology	N/A

14. Adherence to guidelines for child health

Full Name of Indicator	Percentage of children between 0 and 9 years old with complete health check-up according to guidelines
Short name of indicator	Adherence to clinical guidelines for child health
Description	Percentage of children between 0 and 9 years old with exclusive public coverage with complete health check-up according to the guidelines established for each age group.
Comparability	Non-comparable / Country-specific proxy indicator for “Sick child quality score”
VSP Domain and Sub-Domain	Performance / Quality / Provider Competence
Construction	<p><i>Numerator:</i> Number of children between 0 and 9 years old with exclusive public coverage with complete health check-up according to the guidelines</p> <p><i>Denominator:</i> Estimated number of children between 0 and 9 years old with exclusive public coverage during the timeframe</p> <p>Valid cases for this indicator are those that meet the following criteria:</p> <ol style="list-style-type: none"> 1. Child with exclusive public coverage, aged 10 years old or less between the 1st day of the 3rd month and the last day of the measurement interval 2. Compliance with the schedule of specified check-ups 3. All check-ups must submit a weight and height record.
Rationale	Delivery of high-quality care requires the presence of competent providers who provide evidence-based clinical care. This indicator provides information on the extent to which these are observed for the care of adolescents.
Data Source & Year	Administrative data from the Secretaria de Gobierno de Salud, Ministerio de Salud y Desarrollo Social, 2018. For the denominator, data sources are CENSO (2010), EPH y EAHU.
Limitations	Data for this indicator comes from the Programa SUMAR, which provides insurance coverage to selected populations, and covers only public sector facilities.
VSP Methodology	N/A

15. Adherence to guidelines for adolescent health

Full Name of Indicator	Proportion of adolescents with complete health check-ups according to guidelines
Short name of indicator	Adherence to guidelines for adolescent health
Description	Proportion of adolescents between 10-19 years of age with exclusive public coverage who received all health checks according to the protocol established for each age group.
Comparability	Non-comparable / Country-specific proxy indicator for “Adherence to clinical guidelines”
VSP Domain and Sub-Domain	Performance / Quality / Provider Competence
Construction	<p><i>Numerator:</i> Number of adolescents between 10 and 19 years old with exclusive public coverage with complete health check-up according to the guidelines.</p> <p><i>Denominator:</i> Estimated number of adolescents between 10 and 19 years old with exclusive public coverage during the timeframe.</p> <p>Valid cases for this indicator are those that meet the following criteria:</p> <ol style="list-style-type: none"> 1. Adolescents with exclusive public coverage, aged between 10 and 19 years in the considered timeframe 2. Compliance with the schedule of specified controls 3. All controls must present a record of weight, height and blood pressure.
Rationale	Delivery of high-quality care requires the presence of competent providers who provide evidence-based clinical care. This indicator provides information on the extent to which these are observed for the care of adolescents.
Data Source & Year	Administrative data from the Secretaria de Gobierno de Salud, Ministerio de Salud y Desarrollo Social, 2018. For the denominator, data sources are CENSO (2010), EPH y EAHU.
Limitations	Data for this indicator comes from the Programa SUMAR, which provides insurance coverage to selected populations, and covers only public sector facilities.
VSP Methodology	N/A

Safety

No recent indicator available from international or national data sources.

Service Coverage

Coverage looks at the effective application of a broad range of PHC-focused clinical services for the population in need of such services.

RMNCH

16. Demand for family planning satisfied with modern methods

Full Name of Indicator	Demand satisfied with modern methods among women 15-49 years who are married or in a union (%)
Short name of indicator	Demand for family planning satisfied with modern methods
Description	Proportion of married or in-union women of reproductive age (aged 15-49 years) who are married or in a union and have their need for family planning satisfied with modern methods.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / RMNCH
Construction	<p><i>Numerator:</i> Number of married or in-union women of reproductive age (15–49 years old) who are currently using, or whose sexual partner is currently using, at least one modern contraceptive method</p> <p><i>Denominator:</i> Total demand for family planning (the sum of contraceptive prevalence (any method) and the unmet need for family planning)</p>
Rationale	Use of modern contraception is a critical component of women’s, maternal, and population health. This indicator serves as a proxy for population access to reproductive health services, particularly women’s access, which are frequently delivered through the primary health care system and are essential for meeting many health targets. This is SDG indicator 3.7.1.
Data Source & Year	Taken from joint World Bank/WHO “Tracking Universal Health Coverage: 2017 Global Monitoring Report” based on imputed value from regional average. Data are sourced from UNPD estimates based on household surveys, including Demographic and Health Survey (DHS). DHS is a nationally-representative household survey that provides data for a wide range of monitoring and impact evaluation indicators in the areas of population, health, and nutrition. Standard DHS surveys have large sample sizes (usually between 5,000 and 30,000 households) and typically are conducted about every 5 years, to allow comparisons over time.
Limitations	In some surveys, the lack of probing questions, asked to ensure that the respondent understands the meaning of the different contraceptive methods, can result in an underestimation of contraceptive prevalence. Sampling variability may be an issue, particularly when contraceptive prevalence, modern methods is measured for a specific subgroup (according to method, age-group, level of educational attainment, place of residence, etc.) or when analyzing trends over time. This indicator is a measure of both service coverage and fertility preferences and, as such, no target exists. This indicator also specifically addresses only those women who are married or in a union, and may fail to account for any barriers to access encountered by those women who are not but may still desire or benefit from contraception.
VSP Methodology	N/A

17. Antenatal care coverage (4+ visits)

Full Name of Indicator	Antenatal care coverage, four or more visits (ANC4) (%)
Short name of indicator	Antenatal care coverage (4+ visits)
Description	Antenatal care coverage (4+) visits is the percent of women with a live birth who received antenatal care (ANC) 4 or more times.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / RMNCH
Construction	<p><i>Numerator:</i> The number of women aged 15-49 surveyed with a live birth in a given time period who received antenatal care four or more times from any provider</p> <p><i>Denominator:</i> Total number of women aged 15-49 with a live birth in the same period</p>
Rationale	Antenatal care coverage is an indicator of access and use of health care during pregnancy. The antenatal period presents opportunities for reaching pregnant women with interventions that may be vital to their health and wellbeing and that of their infants. Receiving antenatal care at least four times, as recommended by WHO, increases the likelihood of receiving effective maternal health interventions during antenatal visits.
Data Source & Year	Taken from joint World Bank/WHO “Tracking Universal Health Coverage: 2017 Global Monitoring Report” based on data from 2010. Data are sourced from the WHO/RHR global database, which compiles empirical data from DHS, MICS and other national household surveys. Available survey data on this indicator usually do not specify the type of provider; therefore, in general, receipt of care by any provider is measured. At the global level, data from facility reporting are not used. Before data are included into the global databases, UNICEF undertakes a process of data verification that includes correspondence with field offices to clarify any questions regarding estimates.
Limitations	Receiving antenatal care during pregnancy does not guarantee the receipt of interventions that are effective in improving maternal health (effective coverage). Although the indicator for “at least one visit” refers to visits with skilled health providers (doctor, nurse, or midwife), “four or more visits” usually measures visits with any provider because national-level household surveys do not collect provider data for each visit. In addition, standardization of the definition of skilled health personnel is sometimes difficult because of differences in training of health personnel in different countries (UNICEF). Recall error is a potential source of bias in the data.
VSP Methodology	N/A

18. Coverage of DTP3 immunization

Full Name of Indicator	One-year-old children who have received 3 doses of diphtheria-tetanus-pertussis vaccine (DTP3), (%)
Short name of indicator	Coverage of DTP3 immunization
Description	Diphtheria-tetanus-pertussis (DTP) coverage measures the percent of one-year-olds who have received three doses of the combined diphtheria, tetanus toxoid and pertussis vaccine in a given year.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / RMNCH
Construction	<i>Numerator:</i> Number of children of aged 12 months surveyed who have received three doses of the combined diphtheria, tetanus toxoid and pertussis vaccine in a given year <i>Denominator:</i> Total population of children aged 12 months surveyed
Rationale	Immunization is an essential component for reducing under-five mortality. Immunization coverage estimates are used to monitor coverage of immunization services and to guide disease eradication and elimination efforts.
Data Source & Year	Taken from joint World Bank/WHO "Tracking Universal Health Coverage: 2017 Global Monitoring Report" based on data from 2015. The WHO and UNICEF regularly report and release updated immunization coverage data related to the Global Vaccine Action Plan. Data are based on country reported administrative data and household surveys.
Limitations	Given the prevalence of global support for immunization efforts, a high coverage rate of DTP3 immunization may be reflective of strong support from vertical programming in some countries. As such, DTP3 coverage alone is not necessarily a proxy for health system performance.
VSP Methodology	N/A

19. Care-seeking for suspected child pneumonia

Full Name of Indicator	Care-seeking behavior for children with suspected pneumonia (%)
Short name of indicator	Care-seeking for suspected child pneumonia
Description	Percentage of children under 5 years of age with suspected pneumonia (cough and difficulty breathing NOT due to a problem in the chest and a blocked nose) in the two weeks preceding the survey taken to an appropriate health facility or provider.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / RMNCH
Construction	<i>Numerator:</i> Number of children (0-59 months) with suspected pneumonia in the two weeks preceding the survey taken to an appropriate health provider <i>Denominator:</i> Number of children (0-59 months) with suspected pneumonia in the two weeks preceding the survey
Rationale	Pneumonia is a leading cause of child illness and mortality. The strategy for ending preventable child deaths from pneumonia and diarrhea includes a focus on encouraging appropriate care seeking, a key link to receiving appropriate treatment. A number of strategies and programmes to improve care seeking have been developed and implemented in a number of countries.
Data Source & Year	Taken from joint World Bank/WHO "Tracking Universal Health Coverage: 2017 Global Monitoring Report" based on data from 2011. Data are sourced from the UNICEF global database from Demographic and Health Surveys (DHS) and Multiple Indicator Cluster Surveys (MICS).
Limitations	Results are taken from surveys and as a result are subject to recall bias and limitations due to survey design.
VSP Methodology	N/A

20. TB cases detected and treated with success

Full Name of Indicator	Tuberculosis cases detected and treated with success (%)
Short name of indicator	Tuberculosis cases detected and treated with success
Description	Number of new and relapse cases of tuberculosis (TB) that were notified and treated successfully in a given year, divided by the estimated number of incident TB cases in the same year, expressed as a percentage.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / Infectious Diseases
Construction	<i>Numerator:</i> Number of new and relapse cases notified and treated in a given year <i>Denominator:</i> Number of estimated incident cases in the same year
Rationale	This indicator combines case detection rate with treatment success rate to estimate how well the system is detecting and successfully treating TB cases. Treatment success is an indicator of the performance of national TB programs. It also serves as a proxy for a number of aspects of successful service delivery within a health system, including diagnostic and treatment accuracy and the system’s ability to capture and follow up with patients.
Data Source & Year	Taken from joint World Bank/WHO “Tracking Universal Health Coverage: 2017 Global Monitoring Report” based on data from 2014. Estimates of TB incidence are produced through a consultative and analytical process led by WHO and are published annually. These estimates are based on annual case notifications, assessments of the quality and coverage of TB notification data, national surveys of the prevalence of TB disease, and information from death (vital) registration systems. Estimates of incidence for each country are derived, using one or more of the following approaches depending on available data: <ul style="list-style-type: none"> 1. incidence = case notifications/estimated proportion of cases detected; 2. incidence = prevalence/duration of condition; 3. incidence = deaths/proportion of incident cases that die. <p>These estimates of TB incidence are combined with country-reported data on the number of cases detected and treated, and the percentage of cases successfully treated, as described above.</p>
Limitations	The proposed data source for this indicator measures only public sector TB programs and does not include results from private-sector treatment programs or facilities. Therefore, in countries with strong private-sector TB programs, the results do not reflect the totality of the TB treatment success rate.
VSP Methodology	N/A

21. People living with HIV receiving anti-retroviral treatment

Full Name of Indicator	People living with HIV receiving Antiretroviral Therapy (ART) (%)
Short name of indicator	People living with HIV receiving anti-retroviral treatment
Description	Percentage of people living with HIV currently receiving antiretroviral therapy (ART) among the estimated number of adults and children living with HIV.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / Infectious Diseases
Construction	<i>Numerator:</i> Number of adults and children who are currently receiving ART at the end of the reporting period <i>Denominator:</i> Estimated number of adults and children living with HIV
Rationale	ART has been shown to reduce HIV-related morbidity and mortality among people living with HIV and to reduce transmission of HIV. Effective provision of ART can be a marker of how well a health system reaches marginalized populations with higher HIV prevalence.
Data Source & Year	Taken from joint World Bank/WHO "Tracking Universal Health Coverage: 2017 Global Monitoring Report" based on data from 2015. Data are sourced from WHO/UNAIDS estimates. Data on receipt of ART can be collected from facility-based ART registers or drug supply management systems. To estimate the denominator, a standard modelling HIV estimation method, such as in the Spectrum model, is recommended.
Limitations	The indicator permits monitoring trends in coverage but does not attempt to distinguish between different forms of antiretroviral therapy or to measure the cost, quality or effectiveness of, or adherence to the treatment regimen provided. These will each vary within and between countries and are liable to change over time. The indicator measures the number of people provided with medication but does not measure whether the individual took the medication thus it is not a measure of adherence.
VSP Methodology	N/A

22. Children under 5 with diarrhea receiving ORS

Full Name of Indicator	Treatment of diarrhea: Oral rehydration solution (ORS)
Short name of indicator	Children under 5 with diarrhea receiving ORS
Description	The percent of children with diarrhea, a leading cause of death in children under five, who received appropriate treatment with oral rehydration solution.
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / Infectious Diseases
Construction	<p><i>Numerator:</i> Number of children under 5 years of age with diarrhoea in the two weeks preceding the survey given fluid from ORS packets or pre-packaged ORS fluids and zinc supplement</p> <p><i>Denominator:</i> Total number of children aged 0–59 months with diarrhea in the two weeks prior to the survey</p>
Rationale	Diarrhea is a leading cause of child illness and mortality. This is an important indicator of access to health commodities and effective treatment of a common cause of child mortality. This indicator reflects trust in the primary health care system, access to facilities, availability of common home treatments, and health knowledge and behavior.
Data Source & Year	Multiple Indicator Cluster Surveys (MICS), 2011-2012.
Limitations	<p>This indicator does not reflect whether oral rehydration salts and continued feeding were given appropriately. Most diarrhea-related deaths are due to dehydration, and many of these deaths can be prevented with the use of oral rehydration salts at home. However, recommendations for the use of oral rehydration therapy have changed over time based on scientific progress, so it is difficult to accurately compare use rates across countries. Until the current recommended method for home management of diarrhea is adopted and applied in all countries, the data should be used with caution.</p> <p>The prevalence of diarrhea may vary by season. Since country surveys are administered at different times, data comparability is further affected.</p>
VSP Methodology	N/A

Noncommunicable Diseases

23. Prevalence of raised blood pressure (age-standardized estimate)

Full Name of Indicator	Age standardized prevalence of raised blood pressure, regardless of treatment status (%)
Short name of indicator	Prevalence of raised blood pressure (age-standardized estimate)
Description	Age-standardized prevalence of raised blood pressure among persons aged 18+ years (defined as systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg).
Comparability	Comparable / Standard Indicator
VSP Domain and Sub-Domain	Performance / Coverage / NCDs
Construction	<i>Numerator:</i> Number of respondents with systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg <i>Denominator:</i> All survey respondents with a valid measurement
Rationale	Hypertension is a leading risk factor for cardiovascular disease. The results for this indicator represent effective coverage for hypertension, a core part of management of NCDs to reduce complications including renal and cardiovascular disease. This indicator represents a proxy for effective health promotion and service coverage.
Data Source & Year	Taken from joint World Bank/WHO "Tracking Universal Health Coverage: 2017 Global Monitoring Report" based on imputed value from regional average. Data are sourced from NCD-RisC/WHO estimates based on household surveys including Demographic and Health Survey (DHS) and STEPS. DHS is a nationally-representative household survey that provides data for a wide range of monitoring and impact evaluation indicators in the areas of population, health, and nutrition. Standard DHS surveys have large sample sizes (usually between 5,000 and 30,000 households) and typically are conducted about every 5 years, to allow comparisons over time. The STEPwise approach to non-communicable disease risk factor surveillance (STEPS) focuses on obtaining core data at each level on the established risk factors that determine the major disease burden. It is based on survey data and may be supplemented by physical and biometric data.
Limitations	The defined adult population age range differs by country. Rates of normal blood pressure are also influenced by a range of determinants beyond health care service delivery, and thus even appropriate and robust provision of PHC clinical services may only have a limited impact on overall population-based prevalence of some NCDs.
VSP Methodology	For calculation of summary scores in the VSP, this variable was transformed by subtracting the value from 100 to determine the prevalence of normal blood pressure. For this reason, this indicator is shown as "percent of the population with normal blood pressure" on the VSP. These estimates were rescaled to provide finer resolution for the index, based on the observed minima across countries. The rescaled indicator = $(X-50)/(100-50)*100$, where X is the prevalence of normal blood pressure.

Equity

Equity in health service delivery and health outcomes is determined through measures that compare coverage, access and outcome measures across different population groups such as education levels, income, or place of residence.

Where comparable data are available, scores for the Equity domain are color-coded based on the difference between values to reflect good (green), medium (yellow), and poor (red). Scores based on data from non-comparable sources are colored gray.

24. Coverage of RMNCH services, by wealth quintile

Full Name of Indicator	Coverage of RMNCH services: difference between the highest and the lowest wealth quintile
Short name of indicator	Coverage of RMNCH services, by wealth quintile
Description	Difference in RMNCH coverage index for households with mothers that have completed secondary level education versus those without secondary level education.
Comparability	Non-comparable / Country-specific proxy indicator for “Coverage of RMNCH services, by mother’s education”
VSP Domain and Sub-Domain	Equity / Coverage
Construction	This indicator is disaggregated by wealth quintile. Mean-of-means of eight RMNCH interventions, including: <ol style="list-style-type: none"> 1. Early pregnancy (women 20-25 who had a live birth before 18 years old); 2. Antenatal care coverage (at least four visits); 3. Content of antenatal care visits; 4. HIV/AIDS testing (with results) during antenatal care ; 5. Exclusive breastfeeding at 6 months ; 6. Antibiotic treatment for suspected pneumonia; 7. Children aged less than five years with diarrhoea receiving oral rehydration therapy and continued feeding; and 8. Children aged less than five years with pneumonia symptoms taken to a health facility - disaggregated by mother's education.
Rationale	Achieving equitable coverage of basic services is a goal of primary health care.
Data Source & Year	Multiple Indicator Cluster Survey (MICS), 2011-2012.
Limitations	Results are taken from surveys and as a result are subject to recall bias and limitations due to survey design.
VSP Methodology	N/A

25. Under-five mortality rate, by wealth quintile of provinces

Full Name of Indicator	Under-five mortality rate: difference between provinces in the highest and lowest wealth quintile
Short name of indicator	Under-five mortality rate, by wealth quintile of provinces
Description	Difference in under 5 mortality rates between residents of provinces in the highest wealth quintile (with the lowest scores on the Índice de Necesidades Básicas Insatisfechas - NBI) and those of Provinces in the lowest wealth quintile (with the highest scores on the NBI). Probability (expressed as a rate per 1000 live births) of a child born in a specific year or period dying before reaching the age of five years, if subject to age-specific mortality rates of that period.
Comparability	Non-comparable / Country-specific proxy indicator for “Under-five mortality rate, by residence”
VSP Domain and Sub-Domain	Equity / Mortality
Construction	This indicator is disaggregated by wealth quintile of provinces. <i>Numerator:</i> Deaths among children aged 0—4 years (0—59 months of age) <i>Denominator:</i> Number of live births (expressed per 1,000 live births)
Rationale	Achieving equitable health outcomes, across geographic areas, is an essential goal of primary health care. Under-five mortality includes infant and neonatal deaths and reflects the effectiveness of numerous essential services that children receive during their first years of life through primary health care systems, including but not limited to vaccinations, breastfeeding promotion, and nutrition counselling for mothers. It also reflects the social, economic and environmental conditions in which children (and others in society) live. Because data on the incidence and prevalence of diseases (morbidity data) frequently are unavailable, mortality rates are often used to identify vulnerable populations. This indicator captures more than 90% of global mortality among children under age 18.
Data Source & Year	DEIS - Dirección de Estadística e Información de Salud, 2016.
Limitations	The reliability of estimates of under-five mortality depends on the accuracy and completeness of reporting and recording of births and deaths. Underreporting and misclassification are common.
VSP Methodology	N/A